



TITLE:

LONG-RANGE SCATTERING FOR THREE-BODY STARK HAMILTONIANS(Spectrum, Scattering and Related Topics)

AUTHOR(S):

ADACHI, TADAYOSHI

CITATION:

ADACHI, TADAYOSHI. LONG-RANGE SCATTERING FOR THREE-BODY STARK
HAMILTONIANS(Spectrum, Scattering and Related Topics). 数理解析研究所講究録 1994,
873: 120-124

ISSUE DATE:

1994-05

URL:

<http://hdl.handle.net/2433/84090>

RIGHT:

LONG-RANGE SCATTERING FOR THREE-BODY STARK HAMILTONIANS

東大数理科学 足立匡義 (TADAYOSHI ADACHI)

1. INTRODUCTION

Suppose that a system of three particles interacting on each other via the pair potentials $V_{jk}(r_j - r_k)$, $1 \leq j < k \leq 3$, is put in a constant electric field $\mathcal{E} \in \mathbf{R}^d$, $\mathcal{E} \neq 0$. If the mass and charge of j -th particle, $j = 1, 2, 3$, are m_j and e_j and we denote its position vector by $r_j \in \mathbf{R}^d$, the motion of the system is governed by the Hamiltonian

$$\tilde{H} = - \sum_{j=1}^3 \left(\frac{1}{2m_j} \Delta_{r_j} + e_j \langle \mathcal{E}, r_j \rangle \right) + \sum_{1 \leq j < k \leq 3} V_{jk}(r_j - r_k), \quad (1.1)$$

where $\langle x, y \rangle = \sum_{j=1}^d x_j y_j$ for $x, y \in \mathbf{R}^d$.

The purpose of this article is to study the scattering theory for the system in the case when V_{jk} is long-range: $|V_{jk}(y)| = O(|y|^{-\rho})$ as $|y| \rightarrow \infty$ for some $0 < \rho \leq \frac{1}{2}$. To state our result, we need introduce some notations. First we remove the center of mass motion from \tilde{H} . This is achieved as follows: We introduce the metric $r \cdot \tilde{r} = \sum_{j=1}^3 m_j \langle r_j, \tilde{r}_j \rangle$ for $r = (r_1, r_2, r_3)$ and $\tilde{r} = (\tilde{r}_1, \tilde{r}_2, \tilde{r}_3) \in \mathbf{R}^d$ and define $X = \{r \in \mathbf{R}^{3d} \mid \sum_{j=1}^3 m_j r_j = 0\}$ and $X^\perp = \mathbf{R}^{3d} \ominus X = \{r \in \mathbf{R}^{3d} \mid r_1 = r_2 = r_3\}$. Then $L^2(\mathbf{R}^{3d})$ is decomposed into $L^2(X) \otimes L^2(X^\perp)$ and, accordingly, \tilde{H} can be written in the form

$$\tilde{H} = H \otimes Id + Id \otimes T^\perp.$$

Here

$$H = -\frac{1}{2} \Delta - E \cdot x + \sum_{1 \leq j < k \leq 3} V_{jk}(r_j - r_k), \quad T^\perp = -\frac{1}{2} \Delta^\perp - E^\perp \cdot x^\perp, \quad (1.2)$$

Δ (resp. Δ^\perp) is the Laplace-Beltrami operator on X (resp. X^\perp), x and E (resp. x^\perp and E^\perp) are the projections of r and $(\frac{e_1}{m_1} \mathcal{E}, \frac{e_2}{m_2} \mathcal{E}, \frac{e_3}{m_3} \mathcal{E}) \in \mathbf{R}^{3d}$ to X (resp. X^\perp), respectively. We write $V = \sum V_{jk}(r_j - r_k)$. If $E \neq 0$, H is called a three-body Stark Hamiltonian and $H_0 = -\frac{1}{2} \Delta - E \cdot x$ the free Stark Hamiltonian. We are concerned with H and H_0 only.

For each pair $\alpha = (j, k)$, we denote by X^α the configuration space for the relative motion of j -th and k -th particles: $X^\alpha = \{r \in X | m_j r_j + m_k r_k = 0\}$ and $X_\alpha = X \ominus X^\alpha = \{r \in X | r_j = r_k\}$ is the configuration space for the motion of the third particle relative to the center of mass of the pair α . We denote by x^α (resp. x_α) the projection of $x \in X$ to X^α (resp. X_α) and write $V_\alpha(x^\alpha) = V_{jk}(r_j - r_k)$.

We say that $V = \sum_\alpha V_\alpha$ satisfies the condition $(V)_{\rho, \mu}$ if, for each pair α , V_α is a real-valued C^∞ function and satisfies the following conditions:

$$(V.1) \quad |\partial_{x^\alpha}^m V_\alpha(x^\alpha)| \leq C_m \langle x^\alpha \rangle^{-\rho - \mu |m|} \text{ for some } \rho > 0 \text{ and } \mu > 0,$$

$$(V.2) \quad \sum_\alpha \frac{1 + |\omega_\alpha|}{|\omega_\alpha|^2} \sup_{x^\alpha \in X^\alpha} |\omega^\alpha \cdot (\nabla^\alpha V_\alpha)(x^\alpha)| < |E|,$$

where m is any multi-index, $\omega = \frac{E}{|E|}$, $\langle x \rangle = (1 + x^2)^{\frac{1}{2}}$ and ∇^α denotes the gradient on X^α . When V satisfies $(V)_{\rho, \mu}$, both H_0 and H are essentially self-adjoint on $C_0^\infty(X)$, and we denote their closures by the same notations.

The main result of this article is the following theorem.

Theorem 1.1 (Asymptotic Completeness). *Suppose that $E^\alpha \neq 0$ for all pair α and V satisfies the condition $(V)_{\rho, \mu}$ with $0 < \rho \leq \frac{1}{2}$ and $\rho + \mu > 1$. Then the modified wave operators*

$$W_0^{G, \pm} = s - \lim_{t \rightarrow \pm\infty} e^{itH} e^{-\{itH_0 + i \int_0^t V(\frac{E}{2}\tau^2) d\tau\}} \quad (1.3)$$

exist and are unitary operators on $L^2(X)$. Moreover, $W_0^{G, \pm}$ have the intertwining property

$$e^{itH} W_0^{G, \pm} = W_0^{G, \pm} e^{itH_0}, \quad t \in \mathbf{R}. \quad (1.4)$$

Remark 1.2. For long-range potentials, it is known that in general, the usual wave operators

$$W_0^\pm = s - \lim_{t \rightarrow \pm\infty} e^{itH} e^{-itH_0}$$

does not exist even for two-body case (see Ozawa [O]).

Remark 1.3. The condition that $E^\alpha \neq 0$ for all pair α means that the electric field \mathcal{E} is effective on each pair separately, which leads to the unitarity of $W_0^{G, \pm}$. N -body systems with different $\frac{e_j}{m_j}$, $j = 1, \dots, N$, can be treated by our method and the modified wave operators $W_0^{G, \pm}$ for such systems exist and are unitary under the additional assumption that V satisfies the condition corresponding to $(V)_{\rho, \mu}$ with $0 < \rho \leq \frac{1}{2}$ and $\rho + \mu > 1$. However, we shall restrict ourselves here to three-body systems for simplicity.

When $E^\alpha = 0$ for some pair α (note that only one E^α can vanish if $E \neq 0$), the Hamiltonian for the pair α

$$H^\alpha = -\frac{1}{2}\Delta^\alpha + V_\alpha(x^\alpha),$$

where Δ^α is the Laplace-Beltrami operator on X^α , can have bound states and the scattering for H may be multi-channel scattering. Such systems are now under investigation.

Remark 1.4. The condition (V.2) that the pair interactions are small compared with $|E|$ is necessary for us to prove the Mourre estimate and one of propagation estimates. We want to eliminate this condition eventually.

Remark 1.5. The superscript G of the modified wave operators $W_0^{G,\pm}$ indicates that the modification is of Graf-type (see Graf [Gr2] and Jensen-Ozawa [JO2]). In [Gr2], Graf proved the existence and the asymptotic completeness of the modified wave operators $W_0^{G,\pm}$ for two-body systems under the condition that $V \in C^1(\mathbf{R}^d)$ and $\nabla V(x) = O(|x|^{-1-\epsilon})$ as $|x| \rightarrow \infty$ for some $\epsilon > 0$. Our assumption $\rho + \mu > 1$ corresponds to this condition.

Remark 1.6. Under the assumption as in Theorem 1.1, we can prove that the Dollard-type modified wave operators

$$W_0^{D,\pm} = s - \lim_{t \rightarrow \pm\infty} e^{itH} U_0^D(t) \quad (1.5)$$

exist and are unitary operators on $L^2(X)$, where $U_0^D(t)$ is the propagator generated by the time-dependent Hamiltonian

$$H_0^D(t) = H_0 + V(pt - \frac{E}{2}t^2), \quad (1.6)$$

i.e. $\{U_0^D(t)\}_{t \in \mathbf{R}}$ is a family of unitary operators such that for $\psi \in \mathcal{D}(H_0)$, $\psi_t := U_0^D(t)\psi$ is a strong solution of $i\frac{\partial \psi_t}{\partial t} = H_0^D(t)\psi_t$, $\psi_0 = \psi$. For two-body case, Jensen and Yajima [JY] proved this when $d = 1$ and $\mu = \frac{1}{2}$. But we need the assumption $\rho + \mu > 1$.

Remark 1.7. For short-range case, when each pair potential V_α satisfies $|V_\alpha(x^\alpha)| = O(|x^\alpha|^{-\rho})$ as $|x^\alpha| \rightarrow \infty$ for some $\rho > \frac{1}{2}$, the multi-channel scattering mentioned in Remark 1.3 has been studied by Korotyaev [Ko] and Tamura [T2] (see Tamura [T3-4] for general N -body case).

Remark 1.8. When $E = 0$, the problem of the existence and the asymptotic completeness of the multi-channel wave operators has been recently solved for short-range potentials as well as for a class of long-range potentials (cf. Sigal-Soffer [SS1], Graf [Gr1], Kitada [Ki1-2], Tamura [T1] and Yafaev [Y] for short-range case, and Enss [E], Dereziński [D1-2], Gérard [G] and Wang [Wa] for long-range case).

2. AN OUTLINE OF THE PROOF OF THEOREM 1.1

In this section, we give an outline of the proof of Theorem 1.1. For brevity, we omit the proof of the existence of the modified wave operators $W_0^{G,\pm}$. When we admit the existence of $W_0^{G,\pm}$, the intertwining property (1.4) can be proved by the standard way.

First we need the following proposition.

Proposition 2.1. *Assume that V satisfies $(V)_{\rho,\mu}$ with $\rho + \mu > 1$. Let $J \subset \mathbf{R}$ be any bounded interval. Then for $\psi \in \text{Ran} E_J(H)$, there exist $\psi_0^\pm \in L^2(X)$ such that*

$$e^{-itH}\psi = U_0^G(t)\psi_0^\pm + o(1) \text{ as } t \rightarrow \pm\infty. \quad (2.1)$$

We should note that this proposition is a key step for the proof of the asymptotic completeness of $W_0^{G,\pm}$. The property stated in this proposition is called asymptotic clustering by some authors ([SS2], [DG] and [Ki2]). The proof is completed by proving some propagation estimates, but we omit it.

Now we shall show the asymptotic completeness of $W_0^{G,+}$ only. For $W_0^{G,-}$, the proof is similar. In virtue of Proposition 2.1 and the existence of the modified wave operator $W_0^{G,+}$, we see that for $\psi \in \text{Ran} E_J(H)$, there exists $\psi_0^+ \in L^2(X)$ such that $\psi = W_0^{G,+}\psi_0^+$, which implies $\text{Ran} E_J(H) \subset \text{Ran} W_0^{G,+}$. If we note $W_0^{G,+}$ is an isometry and $J \subset \mathbf{R}$ is any bounded interval, this implies that $W_0^{G,+}$ is a unitary operator on $L^2(X)$. This completes the proof of Theorem 1.1.

REFERENCES

- [AH] J.E.Avron and I.W.Herbst, *Spectral and scattering theory of Schrödinger operators related to the Stark effect*, Commun. Math. Phys. **52** (1977), 239–254.
- [CFKS] H.Cycon, R.G.Froese, W.Kirsch and B.Simon, *Schrödinger Operators*, Springer-Verlag, 1988.
- [D1] J.Derezinski, *Algebraic approach to the N-body long-range scattering*, Rev. Math. Phys. **3** (1991), 1–62.
- [D2] J.Derezinski, *Asymptotic completeness of long-range N-body quantum systems*, Ann. of Math. **138** (1993), 427–476.
- [DG] J.Derezinski and C.Gérard, *A remark on asymptotic clustering for N-particle quantum systems*, Schrödinger Operators The Quantum Mechanical Many-Body Problem (E.Balslev, ed.), Lecture Notes in Physics 403, Springer-Verlag, Aarhus, Denmark 1991, pp. 73–78.
- [E] V.Enss, *Long-range scattering of two- and three-body systems*, Proceedings of the Conference ‘Equations aux dérivées partielles’ Saint Lean de Monts, Ecole Polytechnique 1989, pp. 1–31.
- [G] C.Gérard, *Asymptotic completeness for 3-particle long-range systems*, Préprint de l’Ecole Polytechnique, 1992.
- [Gr1] G.M.Graf, *Asymptotic completeness for N-body short-range quantum systems: a new proof*, Commun. Math. Phys. **132** (1990), 73–101.
- [Gr2] G.M.Graf, *A remark on long-range Stark scattering*, Helv. Phys. Acta **64** (1991), 1167–1174.
- [H] I.W.Herbst, *Unitary equivalence of Stark effect Hamiltonians*, Math. Z. **155** (1977), 55–70.
- [JMP] A.Jensen, E.Mourre and P.Perry, *Multiple commutator estimates and resolvent smoothness in quantum scattering theory*, Ann. Inst. H. Poincaré Phys. Theor. **41** (1984), 207–225.

- [JO1] A.Jensen and T.Ozawa, *Classical and quantum scattering for Stark Hamiltonians with slowly decaying potentials*, Ann. Inst. H. Poincaré Phys. Theor. **54** (1991), 229–243.
- [JO2] A.Jensen and T.Ozawa, *Existence and non-existence results for wave operators for perturbations of the Laplacian*, Rev. Math. Phys. **5** (1993), 601–629.
- [JY] A.Jensen and K.Yajima, *On the long-range scattering for Stark Hamiltonians*, J. reine angew. Math. **420** (1991), 179–193.
- [Ki1] H.Kitada, *Asymptotic completeness of N -body wave operators I. Short-range quantum systems*, Rev. Math. Phys. **3** (1991), 101–124.
- [Ki2] H.Kitada, *Asymptotic completeness of N -body wave operators II. A new proof for the short-range case and the asymptotic clustering for the long-range systems*, Functional Analysis and Related Topics, 1991 Proceedings, Kyoto 1991 (H.Komatsu, ed.), Lecture Notes in Mathematics 1540, Springer-Verlag, 1993, pp. 149–189.
- [Ko] E.Korotyaev, *On the scattering theory of several particles in an external electric field*, Math. USSR Sb. **60** (1988), 177–196.
- [M] E.Mourre, *Absence of singular continuous spectrum for certain self-adjoint operators*, Commun. Math. Phys. **78** (1981), 391–408.
- [O] T.Ozawa, *Non-existence of wave operators for Stark effect Hamiltonians*, Math. Z. **207** (1991), 335–339.
- [P] P.Perry, *Scattering Theory by the Enss Method*, Math. Rep. Vol.1, Harwood Academic, 1983.
- [PSS] P.Perry, I.M.Sigal and B.Simon, *Spectral analysis of N -body Schrödinger operators*, Ann. of Math. **114** (1981), 517–567.
- [RS] M.Reed and B.Simon, *Methods of Modern Mathematical Physics, I–IV*, Academic Press.
- [SS1] I.M.Sigal and A.Soffer, *The N -particle scattering problem: asymptotic completeness for short-range systems*, Ann. of Math. **125** (1987), 35–108.
- [SS2] I.M.Sigal and A.Soffer, *Long-range many body scattering: Asymptotic clustering for Coulomb type potentials*, Invent. Math. **99** (1990), 115–143.
- [Sk] E.Skibsted, *Propagation estimates for N -body Schroedinger Operators*, Commun. Math. Phys. **142** (1991), 67–98.
- [T1] H.Tamura, *Asymptotic completeness for N -body Schrödinger operators with short-range interactions*, Commun. Partial Differ. Eqs. **16** (1991), 1129–1154.
- [T2] H.Tamura, *Spectral and scattering theory for 3-particle Hamiltonian with Stark effect: asymptotic completeness*, Osaka. J. Math. **29** (1990), 135–159.
- [T3] H.Tamura, *Spectral Analysis for N -Particle Systems with Stark Effect: Non-Existence of Bound States and Principle of Limiting Absorption*, Preprint, 1993.
- [T4] H.Tamura, *Scattering theory for N -particle Systems with Stark effect: Asymptotic Completeness*, Preprint, 1993.
- [Wa] X.P.Wang, *On the Three-Body Long-Range Scattering Problems*, Lett. Math. Phys. **25** (1992), 267–276.
- [W1] D.White, *The Stark effect and long-range scattering in two Hilbert spaces*, Indiana Univ. Math. J. **39** (1990), 517–546.
- [W2] D.White, *Modified wave operators and Stark Hamiltonians*, Duke Math. J. **68** (1992), 83–100.
- [Y] D.Yafaev, *Radiation conditions and scattering theory for N -particle Hamiltonians*, Commun. Math. Phys. **154** (1993), 523–554.
- [Ya] K.Yajima, *Spectral and scattering theory for Schrödinger operators with Stark-effect*, J. Fac. Sci. Univ. Tokyo Sec IA **26** (1979), 377–390.